

## SECTION 7.12

---

### **Subject: Food Package for Some-Breastfeeding and Non-Breastfeeding Postpartum Women (Food Package 6)**

**References:** 7 CFR 246.10

**Policy:** Food packages must be prescribed by CPAs, in consultation with the participant or parent/caregiver. The food package must be appropriate to the category of eligibility and must be specifically tailored to address the nutritional needs, circumstances, and preferences of the participant.

**Purpose:** To provide a food package that will best meet the participant's needs and will be used by the participant.

#### **Procedures:**

- **Issuance requirements:**

- Food Package 6 is issued to two categories of participants:
  - Non-breastfeeding postpartum women up to 6 months postpartum;
  - Some-Breastfeeding women up to 6 months postpartum, whose infants receive more formula than allowed in the Mostly-Breastfed Infant food package. This includes women breastfeeding multiple infants who are receiving the Some-Breastfeeding Infant food package.  
For maximum formula allowances, refer to
    - Food Package 1: [Contract Formula Amounts Guide](#)
    - Food Package 2: [Contract Formula Amounts Guide](#)
    - Food Package 3: [Formula Amounts Guide for Medical Formula](#)For maximum formula allowances in ounces: [Formula Amounts Guide: Maximum Amounts by Federal Regulation](#).
- For breastfeeding dyads, the infant's food package will determine mom's food package. For infant food packages, refer to [Mom and Baby WIC Food Packages](#).
- Issue the food package as indicated in the table below.
  - Allowed substitutions are described below.
  - Medical documentation is required for some substitutions. Refer to [Section 7.6, Medical Documentation](#).
- Do not provide a food if the participant cannot consume or will not use the food. Examples include
  - Food allergy
  - Medical condition for which the food is inappropriate
  - Vegan family who will not use the food

- **Milk:** Low-fat 1% and fat-free skim are the only types of milk allowed for women.
  - Participants with reported lactose intolerance must try lactose-free milk before milk substitutes may be considered.
  - Other cow's milk options include acidophilus, evaporated and powdered milk. Provide the lowfat and/or fat free versions if available as a food package selection.
  - Goat milk may be provided upon participant request.
  - More than one type of milk or milk substitute may be issued.
  - 1 pound of cheese may be substituted for milk (see table below).
- **Soy-based beverage** may be provided as a milk substitute without medical documentation to replace some or all of the milk, up to the total maximum milk allowance (16 quarts).
- **Tofu:** up to 4 pounds tofu may be substituted for milk without medical documentation. See table below for substitution rates.
  - Amounts > 4 pounds tofu may be provided in cases of qualifying medical need, up to the total maximum milk allowance. Medical documentation must be obtained prior to issuance using [Exhibit 7-E, Authorization of Milk Substitutes](#).
  - For qualifying conditions, see table below.
  - Medical authorization may be provided **by telephone** to a CPA, when necessary to prevent a delay that would place a participant at increased nutritional risk (determined on case-by-case basis) and until written confirmation is received, which must be within 1 month. Voucher issuance is limited to 1 month with verbal verification. Document the temporary verbal approval in the participant's record.
  - Refer to [Section 7.6, Medical Documentation](#).
- **Some-Breastfeeding after 6 months postpartum:** Some-Breastfeeding women receiving Food Package 6 do not receive a food package after the infant turns 6 months. However, these women remain as active participants and receive breastfeeding support, breastpumps (if needed) and nutrition education.
  - If the family reports food insecurity, determine whether a referral to MAC would be appropriate to best meet participant needs.
  - If the participant chooses to participate in MAC rather than WIC in order to benefit from MAC supplemental foods, terminate the WIC certification if still active.
- **Non-breastfeeding women after 6 months postpartum:** Eligibility for non-breastfeeding women ends at 6 months postpartum.
  - Refer these women to the MAC program for supplemental foods.

*See table on next page.*

## Some-Breastfeeding/Non-Breastfeeding Women up to 6 Months Postpartum Food Package 6

Food Item	Maximum Monthly Allowances	Allowed Substitutions	Substitution Rates and Maximums	Medical Documentation Required: Qualifying Reasons
<b>Juice, Single Variety</b>	96 fluid ounces juice	NA		
<b>Milk, Fluid</b>  <i>Cow and Goat milk</i>  <i>Low-fat 1% or fat-free skim milk</i>	16 quarts milk	Cheese	1 pound cheese / 3 quarts of milk <b>Max: 1 pound</b>	NA
		Acidophilus milk Lowfat 1% or Fat-free	<b>Max: 8 half gallon containers</b>	NA
		Lactose-free milk Lowfat 1% or Fat-free	<b>Max: 8 half gallon containers</b>	NA
		Evaporated milk Lowfat or Fat-free	16 fluid ounces / 32 fluid ounces of fluid milk (1:2 ratio) <b>Max: 16 reconstituted quarts = 21 (12 oz.) cans</b>	NA
		Powdered milk Nonfat	Reconstituted amount equal to fluid milk <b>Max: 16 reconstituted quarts</b>	NA
		Soy-based beverage	1 quart soy-based beverage / 1 quart fluid milk <b>Max: 16 quarts</b>	
		Tofu	1 pound tofu / 1 quart of milk <b>Max: 4 pounds</b>	
		If > 4 pounds: 1 pound / 1 quart milk <b>Max: 16 pounds</b>	<ul style="list-style-type: none"> <li>• Milk allergy</li> <li>• Severe lactose maldigestion (cannot tolerate lactose-free milk)</li> <li>• Vegan /vegetarian diet</li> <li>• Medical condition requiring use of soy-based beverage</li> <li>• Religious or cultural observance or practice</li> </ul>	
<b>Breakfast Cereal</b>	36 ounces cereal	NA		
<b>Eggs</b>	1 dozen eggs	NA		
<b>Fruits and Vegetables</b>	\$ 10.00 Cash-value Voucher	NA		
<b>Whole Grains including bread, tortillas, oatmeal or brown rice</b>	NA			
<b>Peanut Butter or Dry/Canned Beans</b>	18 ounces peanut butter <i>or</i> 1 pound dry beans <i>or</i> 64 ounces canned beans	NA		

- **Homeless package:** To better meet the needs of homeless participants, adjust the food package accordingly.
  - Fluid milk quantities may be issued in quart containers.
  - Powdered and evaporated milk are also available.
  - Juice quantities may be issued in the small juice box containers.
  - See [Section 7.14, Homeless Food Package](#).

## Guidance:

- **Lactose intolerance:**
  - Assess the degree of the intolerance. Often lactose intolerance does not require complete avoidance of dairy foods. Determine what approaches the participant has used to reduce symptoms. Individualize counseling. Some approaches include:
    - Introduce milk slowly; gradually increase the amount over time.
    - Drink only small amounts of milk at a time; spread milk intake out during the day (rather than drinking a large amount at one time).
    - Drink milk with food. This will help slow digestion and release of lactose.
    - Yogurt with active and live cultures may help digest the lactose.
    - Try lactose-free milk.
    - Consider some chocolate milk which may be better tolerated.
  - Discuss food package options. Several different types of milk may be provided based on the participant's need.
    - Lactose-free milk
    - Lactose-free milk + cheese
    - Lactose-free milk + regular milk
    - Lactose-free milk + regular milk + cheese
  - Instruct the participant on selecting the lowfat version in the grocery store. It may be helpful to show a picture or sample carton to help her identify the correct item when shopping. When issuing lactose-free milk in HuBERT, select "Fat Free or Lowfat 1%" lactose-free milk.
  - Lactose-free milk may taste different to the participant. Offer suggestions if this is an issue (*note: do not suggest that taste will be an issue*).
    - Make sure the milk is served cold.
    - Use on cereal, mixed with other foods, in hot cocoa, etc. where the difference in taste will not be noticed.
    - If some regular milk is tolerated, mix with lactose-free milk.
    - Try different brands of lactose-free milk.
    - Give it time. Assure the participant that most people become accustomed to the slight difference.

- If the participant expresses reluctance to try lactose-free milk, use open-ended questions to identify and address participant perceptions and concerns.
- **Goat milk** may be preferred by some participants.
  - Goat milk may be useful in some cases of cow milk insensitivity/intolerance or allergy to soy-based beverage. The types of casein in goat milk, plus the evaporation process (for evaporated milk), may render the milk more digestible and less allergenic. The fat has a high proportion of short chain and medium-chain fatty acids and thereby may be more digestible.
  - Goat milk is not appropriate in these situations:
    - Cow's milk protein allergy: cross-reactivity of proteins may increase allergic potential to goat milk.
    - Lactose intolerance: goat milk may not be helpful since the lactose content is comparable to cow's milk.
  - WIC-allowed goat milk is pasteurized and fortified with Vitamin D. Evaporated goat milk is also fortified with folic acid.
- **Peanut butter:**
  - Allergy prevention for the infant: updated AAP guidance states that there is lack of evidence that maternal dietary restrictions during lactation play a significant role in the prevention of allergies in infants. More study is needed to draw any definite conclusions.  
(Source: [\*Effects of Early Nutritional Interventions on the Development of Atopic Disease in Infants and Children\*](#)...PEDIATRICS Volume 121, Number 1, January 2008.)
  - Refer to the Health Care Provider if there are concerns.
- **Soy-based beverage and tofu:**
  - When milk substitutes are used, assess intake and counsel to ensure adequate intake of nutrients that would have been provided by milk. For example:
    - Tofu: does not contain Vitamin D.
    - Tofu often contains less calcium than milk. A 3-ounce tofu serving typically contains 2% to 30% of the Daily Value for calcium, depending on the brand in comparison with 30 % DV for calcium found in 1 cup of milk.