

Multi Year Pregnancy and Postpartum Anemia in Minnesota WIC

Minnesota WIC Information System

	Total Pregnancy ¹	Pregnancy Anemia ²	Total Third Trimester ³	Third Trimester Anemia ⁴	Total Postpartum ⁵	Postpartum Anemia ⁶
MINNESOTA						
2019	17,797	2,894 (16.3)	5,130	1,914 (37.3)	22,462	7,753 (34.5)
2018	18,426	3,469 (18.8)	5,455	2,089 (38.3)	23,972	8,672 (36.2)
2017	20,391	3,152 (15.5)	5,507	2,035 (36.9)	26,295	9,210 (35.0)
2016	21,881	3,232 (14.8)	7,257	2,347 (32.3)	27,847	9,709 (34.9)
2015	21,823	2,932 (13.4)	6,183	2,059 (33.3)	21,952	7,516 (34.2)
2014	23,486	3,052 (13.0)	6,066	1,917 (31.6)	23,844	8,577 (36.0)
2013	24,694	2,682 (10.9)	6,402	1,761 (27.5)	24,518	7,531 (30.7)
2012	25,967	2,642 (10.2)	6,362	1,694 (26.6)	25,796	7,997 (31.0)
2011	24,888	2,447 (9.8)	6,106	1,575 (25.8)	24,709	7,515 (30.4)

**** percentages suppressed if denominator <30

Suggested Citation: Minnesota WIC Information System. *Multi Year Pregnancy and Postpartum Anemia in Minnesota WIC*. Minnesota WIC Program: 2019.

For more information contact: Minnesota WIC Program 1 (800) 657-3942

¹Total number with a valid hemoglobin measurement during pregnancy

²Total number of women during pregnancy with a low hemoglobin (<11.0 mg/dl for first trimester, <10.5 mg/dl for second trimester, and 11.0 mg/dl for third trimester) (women with a valid hemoglobin measurement during pregnancy)

³Total number of women with a valid hemoglobin measurement during the third trimester of pregnancy

⁴Total number of women during the third trimester with a low hemoglobin (<11.0 mg/dl) (women with a valid hemoglobin measurement during the third trimester)

⁵Total number of women with a valid postpartum hemoglobin measurement

⁶Total number of women with low hemoglobin (<12.0 mg/dl) (women with a valid postpartum hemoglobin measurement)