

2025-2026 Combined Respiratory Vaccine Recommendations

| | Influenza Vaccine | RSV Immunization | COVID-19 Vaccine |
|----------------------|---|--|--|
| Infants and Children | All children 6 months and older Some children 6 months to 8 years may need multiple doses AAP, AAFP, CDC | All infants <8 months + children 8-19 months with risk factors (nirsevimab, clesrovimab), typically Oct-Mar, if no maternal RSV vaccine AAP, AAFP, CDC | All children 6-23 months + children 2-18 years with risk factors or if parent requests vaccine AAP, AAFP, IDSA |
| Pregnancy | All At any point in pregnancy ACOG, AAFP, CDC | 32-36 weeks gestation (Pfizer, Abrysvo only) Typically Sept-Jan ACOG, AAFP, CDC | All At any point in pregnancy ACOG, AAFP |
| Adults 19-50 | All AAP, CDC | See Pregnancy AAP, CDC | All Especially if at increased risk for severe infection or never received COVID-19 vaccine AAP, IDSA |
| Adults 50+ | All High-dose, recombinant or adjuvanted flu vaccine preferred for 65+, if available AAP, CDC | All 75+ and adults 50-74 with risk factors One lifetime dose of RSV vaccine AAP, CDC | All Especially for 65+ (two doses), increased risk for severe infection, or never received COVID-19 vaccine AAP, IDSA |

* Either maternal RSV vaccination or infant immunization with RSV monoclonal antibody is recommended.
Most infants will not need both.